VOL. 457 JOURNAL OF CHROMATOGRAPHY DECEMBER 21, 1988

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index)

Effect of random noise and peak asymmetry on the precision and accuracy of measurements of the column efficiency in chromatography
by J. V. H. Schudel and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) (Received August 22nd, 1988) 1

Co-operative cluster model for multivalent affinity interactions involving rigid matrices
by R. J. Yon (London, U.K.) (Received July 26th, 1988) 13

Expert system for the selection of high-performance liquid chromatographic methods in pharmaceutical analysis. Validation of the rules for the selection of the mobile phase
by M. De Smet, A. Peeters, L. Buydens and D. L. Massart (Brussels, Belgium) (Received August 31st, 1988) 25

Numerical simulation of column performance in ion-exclusion chromatography
by B. K. Głód, A. Piasecki and J. Stafiej (Warsaw, Poland) (Received September 1st, 1988) 43

Solubility parameters of gas chromatographic mixed stationary phases
by E. Fernández-Sánchez, A. Fernández-Torres, J. A. García-Domínguez, J. M. Santiuste and E. Pertierra-Rimada (Madrid, Spain) (Received September 1st, 1988) 55

Influence of experimental conditions upon polarity parameters as measured by gas chromatography
by A. Voelkel (Poznań, Poland) (Received August 15th, 1988) 73

Retention in reversed-phase liquid chromatography: solvatochromic investigation of homologous alcohol–water binary mobile phases
by J. J. Michels and J. G. Dorsey (Gainesville, FL, U.S.A.) (Received August 12th, 1988) 85

Reversed-phase high-performance thin-layer chromatography and column liquid chromatography of metal complexes of pheophorbide *a*
by K. Adachi, K. Saitoh and N. Suzuki (Miyagi, Japan) (Received August 25th, 1988) . . 99

Separation of hydrophilic thiols using reversed-phase chromatography with trihaloacetate buffers
by W. A. MacCrehan and D. Shea (Gaithersburg, MD, U.S.A.) (Received May 17th, 1988) 111

Separation of *cis–cis*, *cis–trans* and *trans–trans* isomers of (±)-atracurium besylate and *cis* and *trans* isomers of its major quaternary decomposition products and related impurity by reversed-phase high-performance liquid chromatography
by U. Nehmer (Sofia, Bulgaria) (Received August 24th, 1988) 127

Automated recycling free fluid isotachophoresis. Principle, instrumentation and first results
by J. E. Sloan, W. Thormann, G. E. Twitty and M. Bier (Tucson, AZ, U.S.A.) (Received August 4th, 1988) 137

Photometric and electron-capture modes in a dual-channel sensor
by Y.-Z. Tang and W. A. Aue (Halifax, Canada) (Received August 18th, 1988) 149

Chromatographic method for studying the hygroscopic qualities of solids
by L. G. Berezkina and V. I. Souhodolova (Moscow, U.S.S.R.) (Received July 13th, 1988) 159

High-performance liquid chromatography of proteins on compressed, non-porous agarose beads. I. Hydrophobic-interaction chromatography
by S. Hjertén and J.-L. Liao (Uppsala, Sweden) (Received July 21st, 1988) 165

(Continued overleaf)

High-performance liquid chromatography of proteins on compressed, non-porous agarose beads. II. Anion-exchange chromatography
by J.-L. Liao and S. Hjertén (Uppsala, Sweden) (Received July 21st, 1988) 175

Synthesis of ATP–polyethylene glycol and ATP–dextran and their use in the purification of phosphoglycerate kinase from spinach chloroplasts using affinity partitioning
by L.-O. Persson and B. Olde (Lund, Sweden) (Received August 24th, 1988) 183

Affinity chromatography with triazine dyes immobilized onto activated non-porous monodisperse silicas
by B. Anspach and K. K. Unger (Mainz, F.R.G.) and J. Davies and M. T. W. Hearn (Clayton, Australia) (Received August 8th, 1988) 195

Elution behaviour of some proteins on fresh, acid- or base-treated Sephacryl S-200 HR
by B.-L. Johansson and J. Gustavsson (Uppsala, Sweden) (Received August 22nd, 1988) . 205

Phosphoric acid-modified amino bonded stationary phase for high-performance liquid chromatographic chemical class separation
by J. W. Haas, III, W. F. Joyce, Y.-J. Shyu and P. C. Uden (Amherst, MA, U.S.A.) (Received August 18th, 1988) 215

Characteristics of an avidin-conjugated column in direct liquid chromatographic resolution of racemic compounds
by T. Miwa and T. Miyakawa (Gifu, Japan) and Y. Miyake (Ibaragi, Japan) (Received July 28th, 1988) 227

High-performance liquid chromatography on chiral packed microbore columns with the 3,5-dinitrobenzoyl derivative of *trans*-1,2-diaminocyclohexane as selector
by F. Gasparrini, D. Misiti, C. Villani and F. La Torre (Rome, Italy) and M. Sinibaldi (Monterotondo Stazione, Italy) (Received July 29th, 1988) 235

Indirect method for the determination of water using headspace gas chromatography
by J. M. Loeper and R. L. Grob (Villanova, PA, U.S.A.) (Received August 30th, 1988) . 247

Determination of N-*n*-propylnorapomorphine in serum and brain tissue by gas chromatography–negative ion chemical ionization mass spectrometry
by T. M. Trainor, P. Vouros, P. Lampen, J. L. Neumeyer (Boston, MA, U.S.A.) and R. J. Baldessarini and N. S. Kula (Belmont, MA, U.S.A.) (Received August 31st, 1988) 257

Perfluoro and chloro amide derivatives of aniline and chloroanilines. A comparison of their formation and gas chromatographic determination by mass selective and electron-capture detectors
by H.-B. Lee (Burlington, Canada) (Received August 19th, 1988) 267

Pentafluorobenzylation of capsaicinoids for gas chromatography with electron-capture detection
by A. M. Krajewska and J. J. Powers (Athens, GA, U.S.A.) (Received August 17th, 1988) 279

Identification of methyl diethanolamine degradation products by gas chromatography and gas chromatography–mass spectrometry
by A. Chakma and A. Meisen (Vancouver, Canada) (Received July 27th, 1988) 287

High-performance liquid chromatography of aliphatic aldehydes by means of post-column extraction with fluorometric detection
by H. Koizumi and Y. Suzuki (Yamanashi, Japan) (Received August 3rd, 1988) 299

Chromatographic separation of the diastereomers of a dihydropyridine-type calcium channel antagonist as the bis-3,5-dinitrobenzoates
by J. R. Kern, D. M. Lokensgard and T. Yang (Palo Alto, CA, U.S.A.) (Received August 22nd, 1988) . . . . . .

Large-scale purification of the chromosomal β-lactamase from *Enterobacter cloacae* P99
by C. R. Goward, G. B. Stevens, P. M. Hammond and M. D. Scawen (Salisbury, U.K.) (Received August 1st, 1988) 317

Determination of molluscicidal sesquiterpene lactones from *Ambrosia maritima* (Compositae)
by I. Slacanin, D. Vargas, A. Marston and K. Hostettmann (Lausanne, Switzerland) (Received August 5th, 1988) 325

Simultaneous determination of nine food additives using high-performance liquid chromatography
by Y. Ikai, H. Oka, N. Kawamura and M. Yamada (Nagoya, Japan) (Received August 29th, 1988) 333

Liquid chromatographic determination of bromide after pre-column derivatization to 4-bromoacetanilide
by K. K. Verma, S. K. Sanghi, A. Jain and D. Gupta Jabalpur, India) (Received July 20th, 1988) 345

Notes

Application of inverse gas chromatography to solid propellants
by Y. Ren and P. Zhu (Beijing, China) (Received July 21st, 1988) 354

Electromigration of carrier-free radionuclides. IX. Protolysis of [^{131}I]iodate in aqueous solutions
by F. Rösch, T. K. Hung, M. Milanov and V. A. Khalkin (Moscow, U.S.S.R.) (Received September 8th, 1988) 362

Retention behaviour of diastereomeric truxillic and truxinic diamides and separation of an enantiomeric pair in high-performance liquid chromatography
by S. Caccamese (Catania, Italy) (Received August 18th, 1988) 366

Industrial applications of chromatography. I. Determination of methanol, *n*-butanol and toluene by direct aqueous injection gas chromatography
by A. Horna (Pardubice-Semtín, Czechoslovakia) (Received August 1st, 1988) 372

Study of the adsorption of methyl red on thermally treated gas–liquid chromatographic packings
by G. Pekov, N. Petsev and R. Obretenova (Sofia, Bulgaria) (Received August 17th, 1988) 377

Diazotized dapsone as a reagent for the detection of cannabinoids on thin-layer chromatographic plates
by B. D. Mali and P. P. Parulekar (Aurangabad, India) (Received September 9th, 1988) . 383

Comparison of sulphuric acid treatment and column chromatographic clean-up procedures for the gas chromatographic determination of organochlorine compounds in some food commodities
by P. P. Singh and R. P. Chawla (Ludhiana, India) (Received August 5th, 1988) 387

Horizontal flow-through coil planet centrifuge equipped with a set of multilayer coils around the column holder. Counter-current chromatography of proteins with a polymer-phase system
by Y. Ito and H. Oka (Bethesda, MD, U.S.A.) (Received October 3rd, 1988) 393

Determination of ethylenethiourea in beverages without sample pretreatment using high-performance liquid chromatography and amperometric detection on a copper electrode
by H. Wang, V. Pacáková and K. Štulík (Prague, Czechoslovakia) (Received September 10th, 1988) 398

Simultaneous determination of niacin and niacinamide in meats by high-performance liquid chromatography
by T. Hamano, Y. Mitsuhashi, N. Aoki, S. Yamamoto and Y. Oji (Kobe, Japan) (Received September 5th, 1988) 403

Determination of chlorine and chlorine dioxide in workplace air by impinger collection and ion-chromatographic analysis
by E. Björkholm, A. Hultman and J. Rudling (Solna, Sweden) (Received September 7th, 1988) 409

Non-suppressed ion chromatography of arsenic anions using sodium nitrite solutions as eluents
by N. Hirayama and T. Kuwamoto (Kyoto, Japan) (Received September 9th, 1988) . . . 415

(Continued overleaf)

Contents (continued)

Peak identification of amino acids in liquid chromatography by optical activity detection
by K. C. Chan and E. S. Yeung (Ames, IA, U.S.A.) (Received October 10th, 1988) . . . 421

Determination of terpenic compounds in the essential oil from *Satureja thymbra* L. growing in Sardinia
by W. Capone, C. Mascia, M. Melis and L. Spanedda (Cagliari, Italy) (Received August 9th, 1988) . . . 427

Preparative separation by high-performance liquid chromatography of an extract of oak wood and determination of the composition of each fraction
by J.-L. Puech, P. Rabier and M. Moutounet (Montpellier, France) (Received September 13th, 1988) . . . 431

High-performance liquid chromatographic determination of carbaryl in fruit juices
by R. J. Bushway (Orono, ME, U.S.A.) (Received September 13th, 1988) . . . 437

Determination of isoquinoline alkaloids from Peumus boldus by high-performance liquid chromatography
by P. Pietta and P. Mauri (Milan, Italy) and E. Manera and P. Ceva (Mede, Italy) (Received August 12th, 1988) . . . 442

Book Review

Petroanalysis '87 — Developments in analytical chemistry in the petroleum industry (Proceedings of the Third Petro Analysis Symposium, London) (edited by G. B. Crump) . . . 446

Author Index . . . 447

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)